














# Overall Survival With Circulating Tumor Cell Count–Driven Choice of Therapy in Advanced Breast Cancer: A Randomized Trial

François-Clément Bidard, MD, PhD<sup>1,2</sup> ; Nicolas Kiavue, MD<sup>1</sup> ; William Jacot, MD, PhD<sup>3</sup> ; Thomas Bachelot, MD, PhD<sup>4</sup> ; Sylvain Dureau, PharmD<sup>5</sup>; Hugues Bourgeois, MD<sup>6</sup> ; Anthony Goncalves, MD, PhD<sup>7</sup> ; Etienne Brain, MD, PhD<sup>1</sup> ; Sylvain Ladoire, MD, PhD<sup>8</sup> ; Florence Dalenc, MD, PhD<sup>9</sup>; Joseph Gligorov, MD, PhD<sup>10</sup> ; Luis Teixeira, MD, PhD<sup>11</sup> ; George Emile, MD<sup>12</sup>; Jean-Marc Ferrero, MD, PhD<sup>13</sup>; Delphine Loirat, MD, PhD<sup>1</sup>; Luc Cabel, MD, PhD<sup>1,2</sup> ; Amir Kadi, PhD<sup>5</sup> ; Véronique Diéras, MD<sup>14</sup> ; Catherine Alix-Panabières, PhD<sup>15,16,17</sup>; and Jean-Yves Pierga, MD, PhD<sup>1,17,18</sup>

DOI <https://doi.org/10.1200/JCO.23.00456>

## ABSTRACT

*Clinical trials frequently include multiple end points that mature at different times. The initial report, typically based on the primary end point, may be published when key planned co-primary or secondary analyses are not yet available. Clinical Trial Updates provide an opportunity to disseminate additional results from studies, published in JCO or elsewhere, for which the primary end point has already been reported.*

In patients with hormone receptor–positive, human epidermal growth factor receptor 2–negative advanced breast cancer, the STIC CTC trial established that, for choosing between endocrine therapy (ET) or chemotherapy, the use of circulating tumor cell (CTC) count is noninferior to the investigator’s choice in terms of progression-free survival. Here, we report overall survival (OS) results, a secondary end point. Patients were randomly assigned in a 1:1 ratio to have their first-line treatment (ET or chemotherapy) determined by investigators or CTC count (chemotherapy if  $\geq 5$  CTCs/7.5 mL; ET if low CTC count; CellSearch). OS was assessed at the discontinuation of follow-up. After a median follow-up of 4.7 years, 382 deaths (50.6%) had occurred among 755 patients. Median OS was 51.3 months (95% CI, 46.8 to 55.1) in the CTC arm and 45.5 months (95% CI, 40.9 to 51.1) in the standard arm (hazard ratio [HR] for death, 0.85; 95% CI, 0.69 to 1.03;  $P = .11$ ). Among 189 patients (25.0%) with ET recommended by clinicians and high CTC count, chemotherapy was superior to ET (HR for death, 0.53; 95% CI, 0.36 to 0.78;  $P = .001$ ). In case of a discordant estimate, OS data demonstrate the clinical utility of CTC count.

## ACCOMPANYING CONTENT

-  [Data Sharing Statement](#)
-  [Data Supplement](#)
-  [Protocol](#)

Accepted September 6, 2023

Published November 6, 2023

J Clin Oncol 00:1-7

© 2023 by American Society of Clinical Oncology



[View Online Article](#)

## INTRODUCTION

In hormone receptor–positive metastatic breast cancer (MBC), endocrine therapy (ET) is the preferred option for first-line therapy, but in most international guidelines,<sup>1,2</sup> chemotherapy remains an option for patients with a rapidly evolving disease, the choice between ET and chemotherapy being eventually left to the clinician. In an effort to rationalize, standardize, and optimize the use of frontline chemotherapy, we designed and ran the STIC CTC study, a multicenter randomized phase III trial. This trial compared a choice of first-line therapy (ET or chemotherapy) on the basis of the circulating tumor cell (CTC) count in the experimental arm (CTC arm), with the investigator’s choice in the standard arm. CTC count is a well-established independent prognostic factor in MBC,<sup>3–5</sup> but, to our knowledge, its clinical utility in terms of both progression-free survival (PFS) and overall survival (OS) has never been demonstrated.<sup>3</sup>

The study reached its primary end point, demonstrating that the CTC arm was noninferior to the clinician-driven arm in terms of 2-year PFS.<sup>6</sup> PFS was not significantly superior in the CTC arm, but analyses in the subgroups of patients with discordant Clinical/CTC features suggested the relevance of using the CTC count to choose between ET and chemotherapy. Here, we report the results of updated PFS and final OS (a secondary end point) analyses of the STIC CTC trial.

## METHODS

The design of the STIC CTC trial has been published previously.<sup>6</sup> Detailed trial design and statistical analysis methods are available in the Data Supplement (online only). Briefly, women with a hormone receptor–positive, human epidermal growth factor receptor 2–negative MBC were randomly assigned to a clinician-driven choice of first-line therapy, or to a CTC-driven choice. Before random assignment, the preferred choice of investigators was recorded: chemotherapy (patients referred to as Clin<sup>high</sup>) or ET (patients referred to as Clin<sup>low</sup>).

The baseline CTC count was then determined: patients with a CTC count  $\geq 5$  CTCs/7.5 mL were identified as CTC<sup>high</sup> and those with a count  $< 5$  CTCs/7.5 mL as CTC<sup>low</sup>. Patients randomly assigned in the standard arm received chemotherapy if they were Clin<sup>high</sup> or ET if they were Clin<sup>low</sup>, while patients in the CTC arm received chemotherapy if they were CTC<sup>high</sup> or ET if they were CTC<sup>low</sup>.

Results for the primary end point, 2-year PFS, were previously reported.<sup>6</sup> PFS (with no time boundaries), OS, and subgroup analyses in patients with discordant Clin/CTC features were prespecified secondary end points. The final OS analysis was performed after the follow-up of patients was discontinued, on February 24, 2021.

## RESULTS

From February 1, 2012, to July 28, 2016, 778 patients were randomly assigned to the standard arm (387 patients) or the CTC arm (391 patients; flowchart displayed in the Data Supplement). Patients' characteristics were balanced across the trial arms (Table 1).

Among the 755 participants in the per protocol population, 463 (61.3%) had concordant Clin/CTC features and received the same treatment whatever their allocated arm (ET if CTC<sup>low</sup>/Clin<sup>low</sup>, chemotherapy if CTC<sup>high</sup>/Clin<sup>high</sup>). In a first subgroup with discordant features (Clin<sup>low</sup>/CTC<sup>high</sup>), consisting of  $n = 189$  (25.0%) participants, the 99 Clin<sup>low</sup>/CTC<sup>high</sup> patients allocated to the standard arm received ET, whereas the 90 Clin<sup>low</sup>/CTC<sup>high</sup> patients allocated to the CTC arm received upfront chemotherapy. In the other subgroup with discordant, but opposite, Clin<sup>high</sup>/CTC<sup>low</sup> features ( $n = 103$  [13.6%]), 51 were allocated to the standard arm and received chemotherapy, whereas 52 were allocated to the CTC arm and received ET.

This final analysis was performed after 664 PFS events (87.9% maturity) and 382 deaths (50.6%) had occurred: 197 deaths among 378 patients (52.1%) in the standard arm and 185 among 377 patients (49.1%) in the CTC arm. In the overall population, including all patients with either concordant or discrepant clinical/CTC estimate, the 377 patients of the CTC arm had a median PFS of 15.7 months (95% CI, 12.8 to 17.4) and a median OS of 51.3 months (95% CI, 46.8 to 55.1), whereas the 378 patients of the standard arm had a median PFS of 13.8 months (95% CI, 12.1 to 15.9; hazard ratio [HR] for progression or death, 0.94; 95% CI, 0.81 to 1.09) and a median OS of 45.5 months (95% CI, 40.9 to 51.1; HR for death, 0.85; 95% CI, 0.69 to 1.03; Fig 1A and Fig 1B).

We observed a statistically significant interaction for both PFS and OS between treatment arms and the four subgroups defined by clinical and CTC features (Data Supplement).

Patients with discordant Clin<sup>low</sup>/CTC<sup>high</sup> features derived statistically better outcomes from receiving chemotherapy (in the CTC arm) rather than ET (in the standard arm):

median PFS was 15.7 months (95% CI, 12.7 to 23.2) versus 10.0 months (95% CI, 8.2 to 15.4; HR for progression or death, 0.65; 95% CI, 0.48 to 0.87;  $P = .005$ ), and median OS was 51.8 months (95% CI, 43.3.4 to not reached) versus 35.4 months (95% CI, 30.4 to 45.4; HR for death, 0.53; 95% CI, 0.36 to 0.78;  $P = .001$ ), in the CTC arm and in the standard arm, respectively (Fig 2A and Fig 2B).

By contrast, there was no significant difference in outcomes for the other patient subgroup with discordant Clin<sup>high</sup>/CTC<sup>low</sup> features, in which the CTC-driven decision allowed for a de-escalation from chemotherapy to ET in the CTC arm: median PFS was 9.3 months (95% CI, 6.1.8 to 17.2) versus 14.6 months (95% CI, 10.8 to 20.5; HR for progression or death, 1.14; 95% CI, 0.75 to 1.74;  $P = .54$ ), and median OS was 49.4 months (95% CI, 35.4 to 65.4) versus 45.9 months (95% CI, 36.3 to 59.8; HR for death, 0.88; 95% CI, 0.51 to 1.51;  $P = .64$ ), in the CTC arm and in the standard arm, respectively (Fig 2C and Fig 2D).

Several post hoc analyses were performed (Data Supplement): survival in patients with concordant Clin/CTC status, sensitivity analysis of PFS and OS, variables associated with a CTC<sup>high</sup> status, and proportion of patients having received a CDK4/6 inhibitor as second or later line of therapy; their results did not undermine those of the preplanned analyses.

## DISCUSSION

To our knowledge, the STIC CTC trial is the first contemporary trial that investigated the utility of a prognostic biomarker to drive treatment decision and compared ET or chemotherapy in predefined subgroups of patients with MBC. There was no significant OS benefit in the general population—which comprised a core of approximately 60% patients with concordant Clin/CTC estimate. In subgroup analyses, interaction tests showed that patients with discordant Clin<sup>low</sup>/CTC<sup>high</sup> features, which accounted for about a quarter of the general population, benefited from the proposed strategy, with a clinically and statistically significant gain of 16.4 months in median OS. Interestingly, PFS and OS analyses in patients with Clin<sup>high</sup>/CTC<sup>low</sup> features showed no statistically significant superiority of chemotherapy (in the standard arm) over ET (in the CTC arm), suggesting ET should remain the mainstay of treatment in these patients. Other relevant advantages of CTC count as treatment-driving biomarker over other methods are summarized in the Data Supplement. Importantly, although circulating tumor DNA (ctDNA) is often opposed to CTC, it is important to acknowledge that their clinical validity spectrum is not fully overlapping,<sup>8</sup> and, beyond tumor genotyping, ctDNA clinical utility remains to be formally established by randomized trials in MBC.

The most important limitation to the external validity of our trial is that it was run before the approval of CDK4/6 inhibitors which, given in combination with ET, have

TABLE 1. Patients' Characteristics at Baseline

Characteristic	All Patients	Standard Arm					CTC Arm				
		Clin <sub>low</sub> CTC <sub>low</sub>	Clin <sub>low</sub> CTC <sub>high</sub>	Clin <sub>high</sub> CTC <sub>low</sub>	Clin <sub>high</sub> CTC <sub>high</sub>	Total	Clin <sub>low</sub> CTC <sub>low</sub>	Clin <sub>low</sub> CTC <sub>high</sub>	Clin <sub>high</sub> CTC <sub>low</sub>	Clin <sub>high</sub> CTC <sub>high</sub>	Total
No. of patients	755	176	99	51	52	378	187	90	52	48	377
Age, years, median (range)	63 (30-88)	64 (31-83)	63 (36-87)	60 (42-83)	64 (34-83)	63 (31-87)	64 (36-85)	65 (30-87)	56 (32-81)	61 (33-88)	64 (30-88)
No. of CTCs/7.5 mL of blood, median (range)	2 (0-30,000)	0 (0-4)	14 (5-755)	1 (0-4)	34.5 (5-21,000)	2 (0-21,000)	0 (0-4)	17 (5-30,000)	1 (0-4)	30.5 (5-3040)	2 (0-30,000)
PS, No. (%)											
0	388 (53.7)	104 (61.9)	43 (44.8)	32 (68.1)	16 (32.0)	195 (54.0)	108 (60.0)	39 (45.4)	33 (67.4)	13 (28.3)	193 (53.5)
1	282 (39.1)	57 (33.9)	44 (45.8)	13 (27.7)	25 (50.0)	139 (38.5)	68 (37.8)	35 (40.7)	13 (26.5)	27 (58.7)	143 (39.6)
2/3	52 (7.2)	7 (4.2)	9 (9.4)	2 (4.3)	9 (18.0)	27 (7.5)	4 (2.2)	12 (13.9)	3 (6.1)	6 (13.0)	25 (6.9)
Missing <sup>a</sup>	33	8	3	4	2	17	7	4	3	2	16
Menopausal status, No. (%)											
Premenopausal or perimenopausal	86 (12.2)	20 (12.4)	9 (10.3)	9 (18.0)	5 (10.4)	43 (12.4)	17 (9.5)	6 (6.8)	13 (25.5)	7 (15.9)	43 (11.9)
Postmenopausal	587 (82.9)	136 (84.5)	76 (87.4)	36 (72.0)	40 (83.3)	288 (83.2)	152 (84.9)	79 (89.8)	33 (64.7)	35 (79.6)	299 (82.6)
Unknown	35 (4.9)	5 (3.1)	2 (2.3)	5 (10.0)	3 (6.3)	15 (4.3)	10 (5.6)	3 (3.4)	5 (9.8)	2 (4.5)	20 (5.5)
Missing	47	15	12	1	4	32	8	2	1	4	15
Bone metastases only, No. (%)											
No	555 (73.7)	121 (69.1)	62 (63.3)	40 (78.4)	46 (88.5)	269 (71.5)	137 (73.3)	58 (64.4)	47 (90.4)	44 (91.7)	286 (75.9)
Yes	198 (26.3)	54 (30.9)	36 (36.7)	11 (21.6)	6 (11.5)	107 (28.5)	50 (26.7)	32 (35.6)	5 (9.2)	4 (8.3)	91 (24.1)
Missing	2	1	1	0	0	2	0	0	0	0	0
Liver metastases, No. (%)											
No	600 (79.7)	157 (89.7)	76 (77.6)	40 (78.4)	32 (61.5)	305 (81.1)	162 (86.6)	74 (82.2)	34 (65.4)	25 (52.1)	295 (78.3)
Yes	153 (20.3)	18 (10.3)	22 (22.4)	11 (21.6)	20 (38.5)	71 (18.9)	25 (13.4)	16 (17.8)	18 (34.6)	23 (47.9)	82 (21.7)
Missing	2	1	1	0	0	2	0	0	0	0	0
Histology, No. (%)											
IC-NST	557 (74.6)	150 (86.2)	63 (63.6)	45 (88.2)	28 (54.9)	286 (76.3)	147 (79.4)	49 (55.1)	42 (82.3)	33 (70.2)	271 (72.8)
ILC	153 (20.5)	18 (10.3)	30 (30.3)	4 (7.9)	20 (39.2)	72 (19.2)	29 (15.7)	36 (40.4)	6 (11.8)	10 (21.3)	81 (21.8)
Mixed/others	37 (4.9)	6 (3.5)	6 (6.1)	2 (3.9)	3 (5.9)	17 (4.5)	9 (4.9)	4 (4.5)	3 (5.9)	4 (8.5)	20 (3.4)
Missing	8	2	0	0	1	3	2	1	1	1	5
Estrogen receptor, No. (%)											
Negative	26 (3.4)	7 (4.0)	4 (4.0)	0 (0.0)	2 (3.9)	13 (3.4)	7 (3.7)	2 (2.2)	2 (3.8)	2 (4.2)	13 (3.5)
Positive	681 (90.2)	151 (85.8)	88 (88.9)	49 (96.1)	49 (94.2)	337 (89.2)	169 (90.4)	85 (94.5)	47 (90.4)	43 (89.6)	344 (91.2)
Not done <sup>b</sup>	48 (6.4)	18 (10.2)	7 (7.1)	2 (3.9)	1 (1.9)	28 (7.4)	11 (5.9)	3 (3.3)	3 (5.8)	3 (6.2)	20 (5.3)
Progesterone receptor, No. (%)											
Negative	140 (18.5)	35 (19.9)	18 (18.2)	15 (23.4)	9 (17.3)	77 (20.4)	30 (16.0)	13 (14.4)	9 (17.3)	11 (22.9)	63 (16.7)

(continued on following page)

**TABLE 1.** Patients' Characteristics at Baseline (continued)

Characteristic	All Patients	Standard Arm					CTC Arm				
		Clin <sub>low</sub> CTC <sub>low</sub>	Clin <sub>low</sub> CTC <sub>high</sub>	Clin <sub>high</sub> CTC <sub>low</sub>	Clin <sub>high</sub> CTC <sub>high</sub>	Total	Clin <sub>low</sub> CTC <sub>low</sub>	Clin <sub>low</sub> CTC <sub>high</sub>	Clin <sub>high</sub> CTC <sub>low</sub>	Clin <sub>high</sub> CTC <sub>high</sub>	Total
Positive	537 (71.1)	112 (63.6)	70 (70.7)	32 (62.8)	41 (78.9)	255 (67.5)	139 (74.3)	71 (78.9)	40 (76.9)	32 (66.7)	282 (74.8)
Not done <sup>b</sup>	78 (10.4)	29 (16.5)	11 (11.1)	4 (7.8)	2 (3.8)	46 (12.1)	18 (9.7)	6 (6.7)	3 (5.8)	5 (10.4)	32 (8.5)
Stage at diagnosis, No. (%)											
Synchronous metastasis <sup>c</sup>	45 (26.9)	39 (22.2)	24 (24.2)	8 (15.7)	25 (48.1)	96 (25.4)	56 (30.0)	21 (23.3)	18 (34.6)	12 (25.0)	107 (28.4)
Metachronous metastasis	552 (73.1)	137 (77.8)	75 (75.8)	43 (84.3)	27 (51.9)	282 (74.6)	131 (70.0)	69 (76.7)	34 (65.4)	36 (75.0)	270 (71.6)
Endocrine resistance, <sup>d</sup> No. (%)											
Endocrine-sensitive	539 (71.4)	128 (72.7)	76 (76.8)	27 (52.9)	37 (71.2)	268 (70.9)	141 (75.4)	68 (75.6)	29 (55.8)	33 (68.7)	271 (71.9)
Secondary endocrine resistance	199 (26.4)	45 (25.6)	22 (22.2)	21 (41.2)	15 (28.8)	103 (27.3)	43 (23.0)	21 (23.3)	19 (36.5)	13 (27.1)	96 (25.5)
Primary endocrine resistance	17 (2.2)	3 (1.7)	1 (1.0)	3 (5.9)	0 (0.0)	7 (1.8)	3 (1.6)	1 (1.1)	4 (7.7)	2 (4.2)	10 (2.6)


Abbreviations: CTC, circulating tumor cell; IC-NST, invasive carcinoma of no special type; ILC, invasive lobular carcinoma; PS, performance status.

<sup>a</sup>The information of whether patients were PS 0-1 or 2-3 was available for all participants as it was a stratification factor; however, the exact PS (0 or 1) was not available for all patients.


<sup>b</sup>Data from the latest tumor tissue sampling (for a few patients with a metastasis biopsy, estrogen receptor and progesterone receptor were not assessable, yet all these patients had an estrogen receptor–positive and/or progesterone receptor–positive primary tumor).

<sup>c</sup>If metastases had occurred within 6 months of diagnosis, they were considered as synchronous.

<sup>d</sup>Endocrine resistance status was obtained using the interval between the completion of any previous endocrine therapy and the diagnosis of the metastatic disease, following ABC-4 guidelines.<sup>7</sup>



**FIG 1.** Patients' outcome in the whole population. (A) PFS, all patients. <sup>a</sup>Since the study previously reached its primary end point for PFS noninferiority, no *P* value was generated for this updated analysis. (B) OS, all patients. CTC, circulating tumor cell; HR, hazard ratio; mOS, median overall survival; mPFS, median progression-free survival; OS, overall survival; PFS, progression-free survival.



**FIG 2.** Patients' outcome in subgroups with discordant clinical and CTC estimates. (A) PFS, Clin<sub>low</sub>/CTC<sup>high</sup> patients. (B) OS, Clin<sub>low</sub>/CTC<sup>high</sup> patients. (C) PFS, Clin<sup>high</sup>/CTC<sub>low</sub> patients. (D) OS, Clin<sup>high</sup>/CTC<sub>low</sub> patients. CTC, circulating tumor cell; ET, endocrine therapy; HR, hazard ratio; mOS, median overall survival; mPFS, median progression-free survival; OS, overall survival; PFS, progression-free survival.

significantly prolonged PFS and—for ribociclib—OS.<sup>9–11</sup> Because of the significant survival benefit carried out by CDK4/6 inhibitors, we fully acknowledge that our trial results cannot be extrapolated to CDK4/6 inhibitor-naïve patients. However, the dilemma between ET and chemotherapy persists in patients who have received CDK4/6 inhibitor, either as first-line treatment for MBC or in the adjuvant setting. Our results suggest CTC count may be relevant as an aid to choose between the many ET-based treatment options or

chemotherapy. Trials evaluating CTCs in the CDK4/6 inhibitor era are discussed in the Data Supplement.

In conclusion, the STIC trial established the overall safety of using the CTC count as a standalone biomarker, and despite a lack of significant survival benefit in the general population, it showed that patients with a high CTC count and a low clinical risk estimate may derive a significant OS benefit from chemotherapy.

## AFFILIATIONS

<sup>1</sup>Department of Medical Oncology, INSERM CIC 1428, Institut Curie, Paris and Saint-Cloud, France

<sup>2</sup>Université de Versailles Saint-Quentin, Université Paris-Saclay, Saint-Cloud, France

<sup>3</sup>Department of Medical Oncology, Institut du Cancer Montpellier (ICM), Institut de Recherche en Cancérologie de Montpellier (IRCM), INSERM U1194, Montpellier, France

<sup>4</sup>Department of Medical Oncology, Centre Léon Bérard, Lyon, France

<sup>5</sup>Department of Biostatistics, Institut Curie, PSL Research University, Saint-Cloud, France

<sup>6</sup>Department of Medical Oncology, Victor Hugo Clinic, Le Mans, France

<sup>7</sup>Department of Medical Oncology, Institut Paoli-Calmettes, CRCM, Aix-Marseille University, Marseille, France

<sup>8</sup>Department of Medical Oncology, Centre Georges François Leclerc, Dijon, France

<sup>9</sup>Department of Medical Oncology, Institut Claudius Regaud, IUCT-Oncopole, Toulouse, France

<sup>10</sup>Department of Medical Oncology, INSERM U938, Institut Universitaire de Cancérologie AP-HP Sorbonne Université, Paris, France

<sup>11</sup>Department of Medical Oncology, Hôpital Saint-Louis, AP-HP, Paris, France

<sup>12</sup>Department of Medical Oncology, Centre François Baclesse, Caen, France

<sup>13</sup>Department of Medical Oncology, Centre Antoine Lacassagne, University Côte d'Azur, Nice, France

<sup>14</sup>Department of Medical Oncology, Centre Eugène Marquis, Rennes, France

<sup>15</sup>Laboratory of Rare Human Circulating Cells, University Medical Center of Montpellier, University of Montpellier, Montpellier, France

<sup>16</sup>CREEC, MIVEGEC, University of Montpellier, CNRS, IRD, Montpellier, France

<sup>17</sup>European Liquid Biopsy Society (ELBS), Hamburg, Germany

<sup>18</sup>Université Paris Cité, Paris, France

## CORRESPONDING AUTHOR

François-Clément Bidard, MD, PhD, Institut Curie, 35 rue Dailly, 92210 Saint-Cloud, France; e-mail: francois-clement.bidard@curie.fr.

## PRIOR PRESENTATION

Presented in part at the 2022 San Antonio Breast Cancer Symposium, San Antonio, TX, December 6–10, 2022.

## SUPPORT

Supported by Institut Curie, the French National Cancer Institute (INCa; Program de Soutien aux Techniques Innovantes et Coûteuses STIC 2011), and Menarini Silicon Biosystems (Castel Maggiore, Italy).

F.C.B.'s laboratory is supported by the Institut Curie SIRIC2 SIRIC3 (INCa-DGOS-INSERM-ITMO Cancer\_18000).

## CLINICAL TRIAL INFORMATION

[NCT01710605](https://clinicaltrials.gov/ct2/show/study/NCT01710605) (STIC CTC)

## AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO.23.00456>.

## DATA SHARING STATEMENT

A data sharing statement provided by the authors is available with this article at DOI <https://doi.org/10.1200/JCO.23.00456>.

## AUTHOR CONTRIBUTIONS

**Conception and design:** François-Clément Bidard, Jean-Yves Pierga

**Provision of study materials or patients:** William Jacot, Thomas Bachelot, Hugues Bourgeois, Anthony Goncalves, Etienne Brain, Sylvain Ladoire, Joseph Gligorov, Luis Teixeira, Jean-Marc Ferrero, Catherine Alix-Panabières, Jean-Yves Pierga, Delphine Loirat

**Collection and assembly of data:** William Jacot, Thomas Bachelot, Hugues Bourgeois, Anthony Goncalves, Etienne Brain, Joseph Gligorov, Luis Teixeira, George Emile, Jean-Marc Ferrero, Delphine Loirat, Luc Cabel, Amir Kadi, Véronique Diéras, Catherine Alix-Panabières, Jean-Yves Pierga

**Data analysis and interpretation:** François-Clément Bidard, Nicolas Kiavue, William Jacot, Thomas Bachelot, Sylvain Dureau, Anthony Goncalves, Etienne Brain, Sylvain Ladoire, Florence Dalenc, Joseph Gligorov, Delphine Loirat, Véronique Diéras, Jean-Yves Pierga

**Manuscript writing:** All authors

**Final approval of manuscript:** All authors

**Accountable for all aspects of the work:** All authors

## ACKNOWLEDGMENT

The authors thank Jacques Medioni, MD, PhD (Hôpital Européen Georges-Pompidou, Paris, France), and Hervé Naman (Center Azuréen de Cancérologie, Mougins, France) for their contribution to patient enrollment; Bernard Asselain, MD, PhD, Sandrine Baffert, PhD (both formerly at Institut Curie, Paris, France), and Frédérique Berger, MSc (Institut Curie, Saint-Cloud, France) for their assistance in trial design and analysis, respectively; and Aurore Rampanou, MSc (Institut Curie, Paris, France), for her help with CTC analysis; as well as the patients and their families, the investigators, the caregivers, and the entire STIC CTC team.

## REFERENCES

1. National Comprehensive Cancer Network: NCCN Clinical Practice Guidelines in oncology (NCCN Guidelines): Breast cancer V.7.2021. 2019. <https://www.nccn.org>
  2. Cardoso F, Paluch-Shimon S, Senkus E, et al: 5th ESO-ESMO international consensus guidelines for advanced breast cancer (ABC 5). *Ann Oncol* 31:1623-1649, 2020
  3. Vasseur A, Kiavue N, Bidard F, et al: Clinical utility of circulating tumor cells: An update. *Mol Oncol* 15:1647-1666, 2021
  4. Cristofanilli M, Budd GT, Ellis MJ, et al: Circulating tumor cells, disease progression, and survival in metastatic breast cancer. *N Engl J Med* 351:781-791, 2004
  5. Bidard F-C, Peeters DJ, Fehm T, et al: Clinical validity of circulating tumour cells in patients with metastatic breast cancer: A pooled analysis of individual patient data. *Lancet Oncol* 15:406-414, 2014
  6. Bidard F-C, Jacot W, Kiavue N, et al: Efficacy of circulating tumor cell count–driven vs clinician-driven first-line therapy choice in hormone receptor–positive, ERBB2-negative metastatic breast cancer: The STIC CTC randomized clinical trial. *JAMA Oncol* 7:34-41, 2021
  7. Cardoso F, Senkus E, Costa A, et al: 4th ESO-ESMO International Consensus Guidelines for Advanced Breast Cancer (ABC 4). *Ann Oncol* 29:1634-1657, 2018
  8. Gerratana L, Davis AA, Zhang Q, et al: Longitudinal dynamics of circulating tumor cells and circulating tumor DNA for treatment monitoring in metastatic breast cancer. *JCO Precis Oncol* 5: 943-952, 2021
  9. Hortobagyi GN, Stemmer SM, Burris HA, et al: Overall survival with ribociclib plus letrozole in advanced breast cancer. *N Engl J Med* 386:942-950, 2022
  10. Finn RS, Martin M, Rugo HS, et al: Palbociclib and letrozole in advanced breast cancer. *N Engl J Med* 375:1925-1936, 2016
  11. Johnston S, Martin M, Di Leo A, et al: MONARCH 3 final PFS: A randomized study of abemaciclib as initial therapy for advanced breast cancer. *NPJ Breast Cancer* 5:5, 2019
-

## AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

### Overall Survival With Circulating Tumor Cell Count–Driven Choice of Therapy in Advanced Breast Cancer: A Randomized Trial

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to [www.asco.org/rwc](http://www.asco.org/rwc) or [ascopubs.org/jco/authors/author-center](http://ascopubs.org/jco/authors/author-center).

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](http://Open Payments)).

#### François-Clément Bidard

**Consulting or Advisory Role:** Pfizer, AstraZeneca, Lilly, Novartis, Menarini, Sanofi (Inst), GlaxoSmithKline, Rain Oncology, Caris Life Sciences, GE Healthcare, Exact Sciences, Gilead Sciences, SAGA Diagnostics

**Speakers' Bureau:** Pfizer, Novartis, AstraZeneca, Roche, Lilly, Rain Oncology, Daiichi-Sankyo, Menarini Group

**Research Funding:** Novartis (Inst), Pfizer (Inst), Menarini Silicon Biosystems (Inst), Prolynx (Inst), Merck KGaA (Inst), GE Healthcare (Inst), SAGA Diagnostics (Inst), Rain Oncology (Inst)

**Patents, Royalties, Other Intellectual Property:** ESR1 & MSI detection techniques (patents) (Inst)

**Travel, Accommodations, Expenses:** Roche, Pfizer, AstraZeneca, Novartis, Daiichi-Sankyo

#### Nicolas Kiavue

**Other Relationship:** Grunenthal (Inst)

#### William Jacot

**Honoraria:** AstraZeneca, Eisai, BMSi, Lilly, Daiichi Sankyo Europe GmbH, MSD Oncology, Novartis, Pfizer, Seagen, Roche

**Consulting or Advisory Role:** AstraZeneca, Eisai, Lilly, BMSi, Daiichi Sankyo Europe GmbH, MSD Oncology, Novartis, Roche, Pfizer, Seagen, Gilead Sciences

**Research Funding:** AstraZeneca (Inst), AstraZeneca/Daiichi Sankyo (Inst)

**Travel, Accommodations, Expenses:** AstraZeneca, Novartis, Chugai Pharma, Pfizer, Eisai, Pierre Fabre, GlaxoSmithKline, Roche, Lilly, Sanofi

#### Thomas Bachelot

**Consulting or Advisory Role:** Roche, Novartis, Pfizer, Seagen, Daiichi Sankyo/Astra Zeneca, Lilly

**Research Funding:** Roche (Inst), Novartis (Inst), AstraZeneca (Inst), Seagen (Inst), Pfizer (Inst), Daiichi Sankyo/Astra Zeneca (Inst)

**Travel, Accommodations, Expenses:** Roche, Pfizer, AstraZeneca

#### Hugues Bourgeois

**Travel, Accommodations, Expenses:** Cascadian Therapeutics

#### Anthony Goncalves

**Research Funding:** MSD (Inst), Bristol Myers Squibb (Inst), Novartis (Inst), Boehringer Ingelheim (Inst), Roche/Genentech (Inst), AstraZeneca (Inst), Roche (Inst), Sanofi/Aventis (Inst), Daiichi Sankyo/Astra Zeneca (Inst)

#### Etienne Brain

**Honoraria:** Pfizer, Lilly, Incyte

**Consulting or Advisory Role:** Pfizer, Sandoz-Novartis, Daiichi Sankyo/Astra Zeneca, Menarini

**Travel, Accommodations, Expenses:** Pfizer, Novartis

#### Sylvain Ladoire

**Research Funding:** Novartis (Inst), Eisai (Inst)

**Expert Testimony:** Lilly, Pfizer, Novartis, Bristol Myers Squibb, Astellas Pharma, Roche, Ipsen, Janssen Oncology, Sanofi, Seagen, Daiichi Sankyo/Astra Zeneca, Menarini

**Travel, Accommodations, Expenses:** Pfizer, Novartis, AstraZeneca, Sanofi, Astellas Pharma, Janssen Oncology, Ipsen, Bristol Myers Squibb, Seagen, Daiichi Sankyo/Astra Zeneca

#### Florence Dalenc

**Consulting or Advisory Role:** Pfizer, Seagen, Gilead, AstraZeneca, Daiichi, Novartis (Inst)

**Travel, Accommodations, Expenses:** Gilead and Novartis (Inst)

#### Joseph Gligorov

**Honoraria:** Lilly, Pfizer, Novartis, Menarini, Daiichi Sankyo/AstraZeneca, MSD Oncology, Exact Sciences, Eisai, Eva Pharma, Gilead Sciences, Seagen, Roche/Genentech, AstraZeneca

**Consulting or Advisory Role:** AstraZeneca, Daiichi Sankyo/AstraZeneca, Eisai, Exact Sciences, Gilead Sciences, Lilly, MSD Oncology, Novartis, Pfizer, Roche/Genentech, Seattle Genetics/Astellas

**Speakers' Bureau:** AstraZeneca, Daiichi Sankyo/AstraZeneca, Eva Pharma, Exact Sciences, Gilead Sciences, Lilly, MSD Oncology, Novartis, Pfizer, Roche/Genentech, Seattle Genetics/Astellas

**Travel, Accommodations, Expenses:** Gilead Sciences, Daiichi Sankyo/Astra Zeneca, Pfizer, Novartis, Lilly, Seagen

#### Luis Teixeira

**Consulting or Advisory Role:** Lilly, Daiichi Sankyo/AstraZeneca

**Research Funding:** Novartis, Novartis (Inst)

**Travel, Accommodations, Expenses:** Gilead Sciences

#### George Emile

**Honoraria:** Novartis, Gilead Sciences, Seagen

**Consulting or Advisory Role:** Novartis, Gilead Sciences, AstraZeneca/Daiichi Sankyo

**Research Funding:** Roche (Inst), Bayer (Inst), Seagen (Inst), Lilly (Inst), G1 Therapeutics (Inst)

**Travel, Accommodations, Expenses:** Gilead Sciences, AstraZeneca/Daiichi Sankyo, Novartis, Lilly, Pfizer

#### Jean-Marc Ferrero

**Honoraria:** Novartis, Exact Sciences

**Consulting or Advisory Role:** Pfizer, Daiichi, Novartis

**Travel, Accommodations, Expenses:** Pfizer, Gilead Sciences

#### Delphine Loirat

**Honoraria:** MSD, Gilead Sciences, AstraZeneca, Lilly

**Consulting or Advisory Role:** Roche, MSD Oncology, AstraZeneca, Novartis, Pfizer, Immunomedics, Gilead Sciences, 4D Pharma, Lilly

**Travel, Accommodations, Expenses:** Roche, AstraZeneca, MSD, Pfizer, Gilead Sciences



**Amir Kadi**

**Employment:** Roche (I)

**Véronique Diéras**

**Honoraria:** Roche/Genentech, Novartis, Pfizer, Lilly, AstraZeneca, AbbVie/Abbott, MSD Oncology, Daiichi Sankyo, Seagen, Gilead Sciences, Eisai Europe, Pierre Fabre

**Consulting or Advisory Role:** Roche/Genentech, Novartis, Lilly, Pfizer, AstraZeneca, AbbVie/Abbott, MSD Oncology, Daiichi Sankyo Europe GmbH, Seagen, Gilead Sciences, Eisai Europe, Pierre Fabre, Medac, Menarini Group

**Travel, Accommodations, Expenses:** Roche, Novartis, Pfizer, Lilly, AstraZeneca, Daiichi Sankyo Europe GmbH, Seagen, Gilead Sciences, MSD Oncology

**Jean-Yves Pierga**

**Honoraria:** Novartis (Inst), Daiichi Sankyo/Astra Zeneca (Inst), Gilead Sciences (Inst)

**Consulting or Advisory Role:** Roche/Genentech, Novartis, AstraZeneca, Pfizer, Roche/Genentech (Inst), MSD Oncology (Inst), Lilly, Seagen, Eisai Europe (Inst), Exact Sciences, Veracyte, Daiichi Sankyo Europe GmbH, Gilead Sciences, Menarini Group

**Research Funding:** Roche (Inst), Servier (Inst)

**Travel, Accommodations, Expenses:** AstraZeneca, Gilead Sciences, Pfizer, Daiichi Sankyo Europe GmbH

No other potential conflicts of interest were reported.